

# Comparative analysis of long-term effectiveness of vamorolone vs standard of care glucocorticoid (SoC-GC) treatment in boys with DMD

Craig M. McDonald,<sup>1</sup> Anne-Marie Childs,<sup>2</sup> Liesbeth De Waele,<sup>3,4</sup> Ekaterina Gresko,<sup>5</sup> Michela Guglieri,<sup>6</sup> Jana Haberlova,<sup>7</sup> Shabir Hasham,<sup>5</sup> Marina Katsalouli,<sup>8</sup> Mika Leinonen,<sup>5</sup> Aki Linden,<sup>5</sup> Eugenio Mercuri,<sup>9,10</sup> Francesco Muntoni,<sup>11,12</sup> Yoram Nevo,<sup>13,14</sup> Erik H. Niks,<sup>15</sup> Leanne M. Ward<sup>16</sup>

<sup>1</sup>University of California Davis Health, Davis, CA, USA; <sup>2</sup>Leeds Teaching Hospitals NHS Trust, Leeds, UK; <sup>3</sup>University Hospitals Leuven, Child Neurology, Leuven, Belgium; <sup>4</sup>KU Leuven, Department of Development and Regeneration, Leuven, Belgium; <sup>5</sup>Santhera Pharmaceuticals, Pratteln, Switzerland; <sup>6</sup>The John Walton Muscular Dystrophy Research Centre, Newcastle University and Newcastle Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK; <sup>7</sup>Department of Pediatric Neurology, Second Faculty of Medicine, Charles University and Motol University Hospital, Prague, Czech Republic; <sup>8</sup>Pediatric Neurology Department, St. Sophia's Children's Hospital, Athens, Greece; <sup>9</sup>Catholic University of Sacred Heart, Rome, Italy; <sup>10</sup>Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Rome, Italy; <sup>11</sup>Dubowitz Neuromuscular Centre, Great Ormond Street Institute of Child Health, London, UK; <sup>12</sup>NIHR Great Ormond Street Hospital Biomedical Research Centre, London, UK; <sup>13</sup>Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel; <sup>14</sup>Institute of Pediatric Neurology, Schneider Children's Medical Center of Israel, Petach Tikva, Israel; <sup>15</sup>Department of Neurology, Leiden University Medical Center, Leiden, Zuid-Holland, the Netherlands; <sup>16</sup>Children's Hospital of Eastern Ontario, University of Ottawa, Ottawa, ON, Canada  
Correspondence: cmcdonaldresearch@gmail.com

## Background


- Vamorolone is a first-in-class dissociative steroid approved for the treatment of Duchenne muscular dystrophy (DMD), including in the United States, the European Union, and the United Kingdom.<sup>1-3,8</sup>
- To date, the safety and efficacy of vamorolone has been well established in treatment-naïve boys with DMD up to 30 months of treatment in Phase 2 trials.<sup>4-6</sup>

## Objective

- To assess long-term vamorolone effectiveness, via time to loss of ambulation, and anthropometric changes using longitudinal data from patients enrolled in Phase 2 studies who continued to receive treatment in continued access programs up to inclusion in the Phase 4 GUARDIAN trial, compared with historical data for daily classic corticosteroids (CS; deflazacort or prednisone).
  - Additional bone health outcomes are presented at Poster 62 S.
  - See Poster 66 S for more information about GUARDIAN.

## Methods

### Figure 1. Data sources




\*Propensity score matching criteria: age at start of treatment, age at baseline, and geographical location (North America versus Europe). CINRG-DNHS, Cooperative International Neuromuscular Research Group Duchenne Natural History Study; CUP, compassionate use program; EAP, expanded access program; FOR-DMD, Finding the Optimum Regimen for Duchenne Muscular Dystrophy; NPP, named patient program; US, United States.

- Time to loss of ambulation** was calculated from treatment initiation to time of event.
- Due to differences in study designs and data collection, loss of ambulation was defined as the first of the following:
  - Exact date of loss of ambulation defined as:
    - For vamorolone, the date of 'inability to walk independently' assessed by parent or physician.
    - CINRG-DNHS: participant- or caregiver-reported age at continuous wheelchair use, approximated to nearest month and verified by inability to perform 10-meter walk/run test.
    - FOR-DMD did not collect the exact date of ambulation loss.
  - First occurrence of North Star Ambulatory Assessment rating of 'unable to walk'.
  - Inability to complete the 10-meter walk/run test.
- Anthropometric outcomes** were calculated as z-scores using the Centers for Disease Control growth charts for unaffected males aged 2 to 20 years.<sup>9</sup>
  - Slope analyses from treatment start with random-coefficient models and assumed linear trajectories were used to assess inter-group differences.
- All studies were conducted according to good clinical practice.

## Results

### Figure 2. Subject flow from the Phase 2 studies through to the long-term effectiveness and anthropometric analyses



\*A single boy was originally in the US EAP and transferred to the UK continued access program.  
†Propensity score matching used the following criteria: age at start of treatment, age at baseline, and geographical location (North America versus Europe).  
CINRG-DNHS, Cooperative International Neuromuscular Research Group Duchenne Natural History Study; CUP, compassionate use program; EAP, expanded access program; FOR-DMD, Finding the Optimum Regimen for Duchenne Muscular Dystrophy; NPP, named patient program; US, United States.

## Conclusions

- Vamorolone demonstrated similar effectiveness to classic CS up to a median of 5 years.
- Vamorolone data show that growth can be maintained without earlier loss of ambulation.
- Vamorolone dosage adjustment by clinicians was representative of real-world practice, with a mean dosage of 4.5±1.8 mg/kg/d.
- Like classic CS, vamorolone is associated with increased BMI to be monitored and managed according to guidelines.
- The long-term effectiveness of vamorolone will continue to be assessed in the GUARDIAN study for 3 additional years.

## Effectiveness Results

### Baseline characteristics and dose exposure of the time-to-event effectiveness cohorts

Baseline parameters, mean	Vamorolone 2-6 mg/kg (N=81)	Matched total classic CS (N=287)	Treatment status at assessment	Vamorolone 2-6 mg/kg (N=81)	Matched total classic CS (N=287)
Age, years ± SD	5.8 ± 1.0	6.1 ± 1.2	Mean received dosage, mg/kg/d ± SD	4.5 ± 1.8	N/D
Height z-score ± SD	-0.6 ± 1.0	-0.9 ± 0.9	Median treatment duration, years (range)	5.0 (1.0-8.1)	5.7 (1.0-22.5)
Weight z-score ± SD	0.1 ± 1.0	-0.3 ± 1.0			
BMI z-score ± SD	0.8 ± 1.0	0.5 ± 1.0			
TTSTAND, s ± SD	5.7 ± 3.1	6.7 ± 6.3			
NSAA score, n ± SD	20.7 ± 6.0	21.1 ± 5.8			
Region					
Europe, n (%)	49 (60.5)	97 (33.8)			
US, n (%)	32 (39.5)	190 (66.2)			


BMI, body mass index; CS, corticosteroids; N/D, not determined; NSAA, North Star Ambulatory Assessment; SD, standard deviation; TTSTAND, time to stand from supine; US, United States.

## Key Effectiveness Findings

### Figure 3. Kaplan-Meier curves of time to loss of ambulation for vamorolone 2-6 mg/kg/d versus classic CS (A) and vamorolone 4 to <6 mg/kg/d versus classic CS (B)

- Time to loss of ambulation was similar for vamorolone 2-6 mg/kg/d pooled dosages versus pooled classic CS (Figure 3A), and deflazacort and prednisone individually.
- In a subgroup of boys treated with vamorolone predominantly at 4 to <6 mg/kg/d (4-6 mg/kg/d, <80% of time on 6 mg/kg/d), time to loss of ambulation was similar to the classic CS cohort (Figure 3B).

Individuals with no events were censored at the last follow-up time with documented information or not having met the milestone event. \*The 4 to <6 mg/kg/d cohort were treated with 6 mg/kg/d <80% of the treatment period, acting as a proxy dose for vamorolone 4 mg/kg/d. CS, corticosteroid; HR, hazard ratio; KM, Kaplan-Meier; VAM, vamorolone.



## Anthropometric Results


- The mean (SD) dosage of vamorolone in the anthropometric cohort was 4.5 (1.1) mg/kg/d and the median treatment duration was 4.7 years (range: 1.6-8.1). The classic CS cohort had a median treatment duration of 4.1 years (range: 1.5-10.9).

## Key Anthropometric Findings

### Figure 4. Height (A) and BMI (B) z-score evolution over time since treatment initiation

- Height trajectories were maintained on vamorolone but decreased on classic CS, which showed growth stunting (Figure 4A; growth stunting: height z-score <-2.0 standard deviations).
- After 5 years, the mean height difference between vamorolone- and classic-CS-treated boys was estimated at 11.9 cm.
- BMI z-scores increased in both vamorolone and classic CS cohorts (Figure 4B).

BMI, body mass index; CS, corticosteroid; SE, standard error; VAM, vamorolone.



**Study limitations:** data are from an indirect historical comparison of classic CS that included intermittent dosing in a small number of individuals and measures for loss of ambulation varied across studies.

**References:** 1. AGAMREE 40 mg/ml oral suspension. US Prescribing Information version June 2024; 2. AGAMREE 40 mg/ml oral suspension. EU Summary of product characteristics version January 2026; 3. AGAMREE 40 mg/ml oral suspension. UK Summary of product characteristics version March 2025; 4. Guglieri M, et al. JAMA Neurol. 2022;79(10):1005-14; 5. Dang UJ, et al. Neurology. 2024;102:e208112; 6. Mah JK, et al. JAMA Netw Open. 2022;5:e2144178; 7. Guglieri M, et al. JAMA. 2022;327:1456-68; 8. McDonald CM, et al. Lancet. 2018;391: 451-61; 9. Centers for Disease Control and Prevention. CDC Growth Charts Data Files, [https://www.cdc.gov/growthcharts/percentile\\_data\\_files.htm](https://www.cdc.gov/growthcharts/percentile_data_files.htm) (accessed February 2026).

**Acknowledgements:** We would like to thank all participants, individuals with DMD and their families for participating in the research studies. This study is sponsored by Santhera Pharmaceuticals. Medical writing support was provided by Edward Rochford and funded by Santhera Pharmaceuticals.

**\*Vamorolone should be prescribed according to local approval and indication.**

**Disclosures:** CMM no declarations. AMC clinical trials: Santhera Pharmaceuticals, Sarepta Therapeutics, Entrada Therapeutics, PTC Therapeutics, Roche; consultation: Santhera Pharmaceuticals, Italfarmaco, PTC Therapeutics, Roche. LDW grants: Pfizer; clinical trials: Dyne Therapeutics, Entrada Therapeutics, FibroGen, Génethon, Italfarmaco, Eli Lilly, Pfizer, Prosenza, PTC Therapeutics, Reveragen, Santhera Pharmaceuticals, Sarepta Therapeutics, Wave Life Sciences; consultancy/speaker fees: Entrada Therapeutics, Italfarmaco, Pfizer, PTC Therapeutics, Roche, Santhera Pharmaceuticals, Wave Life Sciences. EG, SH, ML, and AL, employees or consultants of Santhera Pharmaceuticals. MG clinical trials/research: Santhera Pharmaceuticals, Edgewise Therapeutics, Pfizer, Italfarmaco, Roche, ReveraGen, Dyne Therapeutics, Sarepta Therapeutics, PTC Therapeutics; consultancy/speaker fees: Santhera Pharmaceuticals, Roche, Santhera Pharmaceuticals, Wave Life Sciences. JH consultation: Roche, Italfarmaco, Santhera Pharmaceuticals, Pfizer, NS Pharma. MK clinical trials: Santhera Pharmaceuticals, NS Pharma, ReveraGen; consultancy/speaker fees: Genesis Pharma, PTC Therapeutics. EM no declarations. FM consultancy/speaker fees: Santhera Pharmaceuticals, Sarepta Therapeutics, Solid Biosciences, Génethon, Dyne Therapeutics, PTC Therapeutics, Italfarmaco, BioMarin Pharmaceutical, Wave Life Sciences. YN clinical trials: Santhera Pharmaceuticals. EHM clinical trials: Santhera Pharmaceuticals. LMW consultancy: Santhera Pharmaceuticals, Catalyst Pharmaceuticals, Quince Therapeutics, Kye Pharma, Keros, Grunthal; clinical trials: Santhera Pharmaceuticals, Edgewise Therapeutics, Hoffman La-Roche.